

GLASS : GLOBAL LAND-ATMOSPHERE SYSTEM STUDY

Jan Polcher¹, Peter Cox²,
Paul Dirmeyer³, Han Dolman⁴,
Hoshin Gupta⁵, Ann Henderson-Sellers⁶,
Paul Houser⁷, Randy Koster⁷, Taikan Oki⁸,
Andy Pitman⁹, and Pedro Viterbo¹⁰

¹Laboratoire de Meteorologie Dynamique du CNRS,
²Hadley Centre, ³Center for Ocean-Land-Atmos. Studies,
⁴Alterra, ⁵University of Arizona, ⁶ANSTO,
⁷NASA/Goddard Space Flight Center,
⁸University of Tokyo, ⁹Macquarie University
¹⁰European Centre for Med.-Range Weather Forecasts

A new generation of land surface schemes (LSS) is emerging. The schemes are evolving from general circulation model parameterizations that provide fluxes to the atmosphere into independent models that are increasingly being compared to models of hydrology, biogeochemistry and ecology. This expanding scope is driven by the growth of interdisciplinary studies of the earth system. GLASS aims to encourage these developments by coordinating the evaluation and intercomparison of this new generation of LSSs, and applying them to scientific queries of broad interest.

Workshops organized by GEWEX, the European Centre for Medium-Range Weather Forecasts (ECMWF), the International Geosphere-Biosphere Programme (IGBP), and the Institut National des Sciences de l'Univers (INSU) during the past few years have pointed out a number of directions in which innovation in the next generation of LSSs will be necessary. The **first point** identified was the inclusion of bio-physical processes. There is a need to **include the carbon cycle** in order to be able to provide the atmospheric models with CO₂ fluxes and to simulate the evolution of the vegetation and thus the feedbacks between climatic variations and the state of the biosphere. The **second feature** that will set this new generation apart will be the **larger importance given to the horizontal complexity of the surface**. Indeed, previous LSSs were characterized by a very sophisticated description of the vertical processes while retaining simple assumptions on the horizontal variance of surface conditions. The **final innovation** to be included in the next generation will be **data assimilation methods** for either the analysis of the state of the surface or the evaluation of surface schemes at the global scale. The remotely sensed variables that will become available in the years to come will demand

from LSSs new methods for taking advantage of the new information. With this increase in complexity, new feedbacks between the atmosphere and the surface will be simulated and a more systematic analysis of the sensitivity of the climate to surface processes is needed.

Land surface scheme validation and intercomparison projects such as the Project for Intercomparison of Land-surface Parameterization Schemes (PILPS) (Henderson-Sellers et al., 1995) and the Global Soil Wetness Project (GSWP) (Dirmeyer et al., 1999) helped to assess the previous generation of LSSs. PILPS made major contributions to their evaluation at the local scale using observed atmospheric forcing and high quality measured fluxes for the validation of the simulated processes. In GSWP on the other hand, the LSSs were used in a mode much closer to their application in GCMs (i.e., on a global grid with a resolution coarser than the typical spatial scales of surface processes). Both of these projects proved very useful and need to be continued with the next generation for the validation of added processes. However, they also need to be supplemented by new experiments.

As land surface schemes are applied from the plot scale to the regional scale and in a forced or coupled mode, the new experiments will specifically have to address the issues of sub-grid scale variability of the surface and the feedbacks existing between surface processes and the atmosphere. The complementarity of the various applications of LSSs provides a richness of research and can be illustrated, for instance, with the case of simulated soil moisture. In the PILPS experiments the soil moisture simulated with the atmospheric forcing observed at a site could be compared to the measured values (Shao and Henderson-Sellers, 1996). In GSWP, only the soil moisture anomalies at 1°x1° could be evaluated as the models use different water holding capacities, which were all different from the one that can be deduced from observed values (Entin et al., 1999). When an LSS is coupled to an atmospheric column model the relationship between the soil moisture and the atmospheric conditions can be explored at the local scale. This set-up was used by Douville et al. (2000) to determine how the assimilation of near surface temperature and humidity could help improve the simulated soil moisture. Finally, with coordinated GCM sensitivity experiments the impact of the different approaches to soil moisture modelling on climate change could be evaluated (Crossley et al., 2000). Unfortunately, we are

not presently able to say how the results obtained in PILPS can be related to the uncertainty in soil moisture changes found for a climate with increased greenhouse gas concentrations. Thus, a coordination of projects is needed to take full advantage of the diversity of application areas for LSSs.

The aim of GLASS is to foster an evaluation of the next generation of LSSs and to coordinate the evaluation of LSSs in their different applications. GLASS will also serve as an interface between the land-surface community and other GEWEX projects. The proposed structure of GLASS (see figure below) highlights the spatial scales at which the schemes are applied and the degree of interaction allowed with the atmosphere. This denotes four actions which will coordinate intercomparisons in their field of land-surface scheme applications. GLASS will also include one transversal action [Assistance for Land-surface Modelling Activities (ALMA)] which will provide an infrastructure and technical support for these intercomparisons.


The action dedicated to local scales and off-line simulations is the continuation of the PILPS project. **The next intercomparisons will be carried out over the Torne basin in Sweden (see article on page 8) and over a mature forest at the Russian Valdai site.** These experiments will allow an evaluation of cold-season processes in LSSs. It is expected that in the near future a PILPS experiment will be launched to evaluate the newly gained ability of land-surface schemes to simulate CO₂ fluxes.

The action concentrating on global off-line applications of LSSs is the continuation of GSWP. Before the next phase of GSWP can be launched with the International Satellite Land Surface Climatology Project (ISLSCP) Initiative II data, sensitivity experiments will be carried out to evaluate the impact of errors in the forcing data. Experiments are also planned to evaluate our ability to take into account the spatial variability of soil wetness at a 1°x1° resolution. Later, with the 10 years of ISLSCP Initiative II data the inter-annual variability of LSSs will be evaluated.

The local coupled action will concentrate on data assimilation issues. The first step towards this goal will be to set up a simplified coupled system in which LSSs and their interaction with the atmosphere can be compared. With this tool in hand the simulated interactions between sub-grid variability and the planetary boundary layer will be compared. This will also serve to evaluate data assimilation methods for land-surface schemes.

In the global coupled action, the sensitivity of the atmosphere to land-surface processes will be studied using an ensemble of general circulation models and LSSs. The two issues that are most in need of model independent assessments are the role of climate variability and the sensitivity of climate to anthropogenic forcing. This action will work in close collaboration with the two Atmospheric Model Intercomparison Project (AMIP) diagnostic sub-projects which deal with the validation of surface processes in GCM simulations.

The infrastructure action, ALMA, will help the four actions described above by providing data exchange standards and software to manage the data.



ALMA will also help participating groups with the implementation of the PILPS-4c coupler (Polcher et al., 1998) and ensure the exchange of software and expertise with the aim of facilitating the participation in GLASS projects.

GLASS will be managed by a science panel composed of the leaders of the five actions and representatives of the GEWEX Hydro-meteorology Panel (GHP), ISLSCP, IGBP/Biospheric Aspects of the Hydrologic Cycle (BAHC) and the Working Group on Numerical Experimentation (WGNE). Its role will be to coordinate the five actions and ensure that all application areas of LSSs benefit from progress made in the various projects. The panel will hold its first meeting this summer and at that time the implementation plan of GLASS will be finalized. The first local scale off-line experiment (PILPS-2e) will be launched this month and the data exchange protocol to be used has been developed in collaboration with ALMA. Other actions have submitted projects for funding and will start before the end of the year. The evolution of GLASS can be followed at its **Web site: <http://hydro.iis.u-tokyo.ac.jp/GLASS/>**.

References

Crossley, J., J. Polcher, P. Cox, N. Gedney, and S. Planton, 2000. Uncertainties linked to land-surface processes in climate change simulations. *Clim. Dyn.*, in press.

Dirmeyer, P. A., A. J. Dolman, and N. Sato, 1999. The global soil wetness project: A pilot project for global land surface modeling and validation. *Bull. Amer. Meteor. Soc.*, 80:851-878.

Douville, H., P. Viterbo, J.-F. Mahfouf, and A. C. M. Beljaars, 2000. Evaluation of the optimum interpolation and nudging techniques for soil moisture analysis using FIFE data. *Mon. Wea. Rev.*, 128:in press.

Entin, J. K., A. Robock, K. Y. Vinnikov, V. Zabelin, S. Liu, A. Namkhai, and T. Adyasuren, 1999. Evaluation of global soil wetness project soil moisture simulations. *J. Meteor. Soc. Japan*, 77:17-32.

Henderson-Sellers, A., A. J. Pitman, P. K. Love, P. Irannejad, and T. Chen, 1995. The project for intercomparison of land-surface parameterization schemes (PILPS): Phase 2 and 3. *Bull. Amer. Meteor. Soc.*, 76:489-503.

Polcher, J., B. McAvaney, P. Viterbo, M.-A. Gaertner, A. Hahmann, J.-F. Mahfouf, J. Noilhan, T. Phillips, A. Pitman, C. A. Schlosser, J.-P. Schulz, B. Timbal, D. Verseghy, and Y. X., 1998. A proposal for a general interface between land-surface schemes and general circulation models. *Glob. and Plan. Change*, 19:263-278.

Shao, Y. and A. Henderson-Sellers, 1996. Modelling soil moisture: a project for intercomparison of land surface parameterization schemes phase 2(b). *J. Geophys. Res.*, 101(D3):7227-7250.